

AD-753 470

EQUIVALENT FORMULATIONS OF THE HIRSCH
CONJECTURE FOR ABSTRACT POLYTOPES

John A. Lawrence, Jr.

California University

Prepared for:

Office of Naval Research
National Science Foundation

November 1972

DISTRIBUTED BY:

NTIS


National Technical Information Service
U. S. DEPARTMENT OF COMMERCE
5285 Port Royal Road, Springfield Va. 22151

ORC 72-28
NOVEMBER 1972

EQUIVALENT FORMULATIONS OF THE HIRSCH CONJECTURE FOR ABSTRACT POLYTOPES

by
JOHN A. LAWRENCE, JR.

AD753470



NATIONAL TECHNICAL
INFORMATION SERVICE
U.S. DEPARTMENT OF COMMERCE
WASHINGTON, D.C. 20540

OPERATIONS
RESEARCH
CENTER

DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited

COLLEGE OF ENGINEERING
UNIVERSITY OF CALIFORNIA • BERKELEY

18

EQUIVALENT FORMULATIONS OF THE HIRSCH CONJECTURE FOR ABSTRACT POLYTOPES

by

John A. Lawrence, Jr.
Operations Research Center
University of California, Berkeley

NOVEMBER 1972

ORC 72-28

This research has been partially supported by the Office of Naval Research under Contract N00014-69-A-0200-1010 and the National Science Foundation under Grant GK-23153 with the University of California. Reproduction in whole or in part is permitted for any purpose of the United States Government.

Unclassified

Security Classification

DOCUMENT CONTROL DATA - R & D

Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified.

1. ORIGINATING AGENCY (Corporate author)		2a. REPORT SECURITY CLASSIFICATION	
University of California, Berkeley		Unclassified	
		2b. GROUP	
3. REPORT TITLE			
EQUIVALENT FORMULATIONS OF THE HIRSCH CONJECTURE FOR ABSTRACT POLYTOPES			
4. DESCRIPTIVE NOTES (Type of report and, inclusive dates)			
Research Report			
5. AUTHOR(S) (First name, middle initial, last name)			
John A. Lawrence, Jr.			
6. REPORT DATE		7a. TOTAL NO. OF PAGES	7b. NO. OF REFS
November 1972		1318	2
8a. CONTRACT OR GRANT NO.		9a. ORIGINATOR'S REPORT NUMBER(S)	
GK-23153		ORC 72-28	
b. PROJECT NO.			
c.		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.			
10. DISTRIBUTION STATEMENT			
This document has been approved for public release and sale; its distribution is unlimited.			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY	
Also supported by the Office of Naval Research under Contract N00014-69-A-0200-1010.		NSF-GK-23153-Shephard The National Science Foundation Washington, D.C. 20550	
13. ABSTRACT			
SEE ABSTRACT.			
II			

ABSTRACT

Abstract polytopes are mathematical creations which are defined by three axioms. It has been shown that simple polytopes are a proper subclass of abstract polytopes. Hence theorems proving facts about abstract polytopes in general, prove facts about simple polytopes in particular.

Klee and Walkup [2] showed the following four statements were mathematically equivalent for simple polytopes:

- i) Any two vertices of a simple polytope can be joined by a W_v (nonreturning) path.
- ii) $\Delta(n,d) \leq n - d$ (Hirsch conjecture).
- iii) $\Delta(2d,d) \leq d$.
- iv) For a Dantzig figure, (P,x,y) , $\delta_p(x,y) = d$.

The purpose of this paper is to show that the four statements above are equivalent for the larger class of abstract polytopes as well. Thus, it is possible to tackle the problem of the well-known Hirsch conjecture by applying the well defined structure and theorems of abstract polytopes to any of the above statements.

1. INTRODUCTION

In Adler [1], it was shown that the class of simple polytopes is a proper subclass of the class of abstract polytopes. Because of the simplicity and mathematical precision which define abstract polytopes, they become interesting and practical tools for investigating properties of simple polytopes.

Klee and Walkup [2], showed that the well known Hirsch conjecture is mathematically equivalent to three other statements. In this paper it will be shown that a corresponding set of statements are equivalent for abstract polytopes, and hence abstract polytope theory may shed new light in proving or disproving the Hirsch conjecture.

The method of proof will be the method of consecutive implication, i.e. for statements, i, ii, iii, iv, it is shown $i \Rightarrow ii \Rightarrow iii \Rightarrow iv \Rightarrow i$. Three implications are trivially proved. A constructive proof for the fourth implication is developed strictly from abstract polytope theory and is based on Klee and Walkup [2].

Finally an example of the construction is presented in the appendix.

2. NOTATIONS AND DEFINITIONS

Let C be a set of distinct symbols.

Any subset v , of d distinct symbols contained in C is called a vertex.

$|v|$ denotes the number of symbols in v .

Vertices u, v of d symbols are said to be neighbors iff $|u \cap v| = d - 1$.

A path from v_0 to v_k is a series of vertices v_0, v_1, \dots, v_k s.t. v_i, v_{i+1} are neighbors, $i = 0, \dots, k-1$.

Definition of an Abstract Polytope

P is a labeled abstract polytope consisting of vertices $V(P)$ iff

(A1) $v \in V(P) \Rightarrow |v| = d$.

(A2) $v \in V(P) \Rightarrow$ for any symbol $T \in v$, \exists exactly one other vertex $w \in V(P)$, s.t. $v - T \subset w$.

(A3) For any $v_0, v_k \in V(P)$, \exists a path $R : v_0, v_1, \dots, v_k$ s.t. $v_0 \cap v_k \subset v_i \forall v_i \in R$. (R is called an A3 path.)

Note: We will use P also to denote the union of all symbols used in the vertices of $V(P)$.

$P(n, d) \equiv$ set of all abstract polytopes

s.t. $|P| = n, |v| = d$.

If $v \in V(P)$ and some set $S \subset v$ such that $|S| = k (> 0)$, S is said to generate a $d - k$ face on P ; i.e., $F_P(S)$, the face generated by S on P , consists of all $v \in V(P)$ s.t. $S \subset v$. It is easily verified that after the k symbols

common to $v \in V(F_p(S))$ have been dropped we have formed a new abstract polytope $Q \in \mathcal{P}(n', d - k)$ where $n' \leq n - k$. We say Q corresponds to $F_p(S)$ and the vertices of Q have a one to one correspondence to the vertices of $F_p(S)$, thus $q \in V(Q)$ corresponds to $q \cup S \in V(F_p(S))$.

$\delta_p(x, y) \equiv$ the *distance* from x to y on P , i.e., the length of the shortest path between x and y .

$$\Delta_a(n, d) \equiv \text{the max diameter of } \mathcal{P}(n, d) \\ \equiv \max_{P \in \mathcal{P}(n, d)} \max_{x, y \in V(P)} \delta_p(x, y).$$

A *nonreturning (NR)* path from v_0, v_1, \dots, v_k on P is a path s.t. the symbol $T \in v_i, T \notin v_{i+1} \Rightarrow T \notin v_j \quad i + 1 \leq j \leq k$.

A *Dantzig figure* (P, x, y) is an abstract polytope $P \in \mathcal{P}(2d, d)$ s.t. \exists vertices $x, y \in V(P)$ s.t. $x \cap y = \emptyset$.

3. THEOREM

The following four statements are equivalent for abstract polytopes:

- i) \exists an NR path between any two vertices $x, y \in V(P)$.
- ii) $\Delta_a(n, d) \leq n - d$.
- iii) $\Delta_a(2d, d) \leq d$.
- iv) For every Dantzig figure (P, x, y) , $\delta_P(x, y) = d$.

Proof: (by consecutive implication)

- (a) $i \Rightarrow ii$ if \exists an NR path between $x, y \in V(P)$ then

$$\delta_P(x, y) \leq n - (|x \cap y| + d). \text{ If } x \cap y = \emptyset \text{ result follows.}$$

Say $x \cap y = S$, $|S| = k$. Consider $F_P(S)$. Then

$$\delta_{F_P(S)}(x, y) \leq n' - (d - k) \leq n - k - (d - k) = n - d.$$

$$\therefore \delta_P(x, y) \leq n - d \quad \forall P \in \mathcal{P}(n, d).$$

- (b) $ii \Rightarrow iii$ trivial. Let $n = 2d$.

- (c) $iii \Rightarrow iv$ $\delta_P(x, y) \leq d$ by iii). But since $x \cap y = \emptyset$, $\delta_P(x, y) \geq d$

$$\therefore \delta_P(x, y) = d.$$

- (d) $iv \Rightarrow i$

Consider $x, y \in V(P)$. Assume $x \cap y = \emptyset$ for convenience. If $x \cap y = S (\neq \emptyset)$ consider $F_P(S)$. After eliminating S from each vertex of $F_P(S)$ we have a new abstract polytope, Q , whose vertices correspond to $F_P(S)$. In particular the vertices which correspond to x and y have an empty intersection. Thus, we can apply the following theory to Q and make the corresponding transformation back to $F_P(S)$ and hence P .

Assume $n > 2d$. (If $n = 2d$ result follows trivially since $\delta_P(x, y) = d$.)

Notation

Let $M = P - (x \cup y)$; say $|M| = m$, i.e., $m = n - 2d$.

Form a set M' of new symbols s.t. $T \in M \Rightarrow T' \in M'$.

Now we form P^* by the algorithm, illustrated in Figure 1.

We now show $P^* \in (n + m, d + m)$. We show the more general fact:

Lemma 1:

$$P^j \in P(n + j, d + j) .$$

Proof: (by induction)

Trivially $P^0 \in P(n, d)$.

Suppose $P^{j-1} \in P(n + j - 1, d + j - 1)$.

Consider P^j . It is easily verified A1 and A2 are satisfied.

To show A3, consider the following three cases. Suppose $u^j, v^j \in V(P^j)$.

Case I: $T' \in u^j, T' \in v^j$

$\exists u^{j-1}, v^{j-1} \in V(P^{j-1})$ s.t. $u^{j-1} = u^j - T', v^{j-1} = v^j - T'$.

Since \exists an A3 path $u^{j-1} = u_0^{j-1}, \dots, u_k^{j-1} = v^{j-1}$ on P^{j-1} .

Let $u_i^j = u_i^{j-1} \cup T'$. This is the required A3 path on P^j .

Case II: $T' \in u^j, T' \notin v^j$

By construction, $T' \notin v^j \Rightarrow T \in v^j$ and $\exists y^j \in V(P^j)$ s.t. $y^j = v^j \cup T - T'$.

By Case I \exists an A3 path, R , on P^j from u^j to y^j . \therefore The required A3 path from u^j to v^j is R amended with the edge (y^j, v^j) .

Case III: $T' \notin u^j, T' \notin v^j$

Let $y^j, z^j \in V(P^j)$ be such that $y^j = u^j \cup T' - T, z^j = v^j \cup T' - T$.

Construction 1

Let $j = 0$, $P^0 = P$, $M^0 = M$, $M'^0 = M'$.




Figure 1

We know \exists an A3 path R from y^j to z^j on P^j . $R: y^j = w_0^j, \dots, w_f^j, \dots, w_\ell^j, \dots, w_k^j = z^j$ where for $f \leq i \leq \ell$, $T \in w_i^j$. (It is not necessary that $T \in w_i^j$ for any i or for only one interval.)

Consider the path $\bar{R}: u^j = \bar{w}_0^j, \dots, \bar{w}_f^j, \dots, \bar{w}_\ell^j, \dots, \bar{w}_k^j = v^j$ s.t. for $0 \leq i \leq f-1$ and $\ell+1 \leq i \leq k$, $\bar{w}_i^j = w_i^j \cup T' - T$ and for $f \leq i \leq \ell$, $\bar{w}_i^j = w_i^j$.

It is not difficult to verify \bar{R} exists and is the required A3 path from u^j to v^j on F^j . ■

Corollary 1:

$P^* \in P(n+m, d+m)$, i.e., $P^* \in (2(n-d), (n-d))$.

We now make the following observations:

Observation 1: \exists nodes $x^*, y^* \in V(P^*)$ s.t. $x^* = x \cup M'$, $y^* = y \cup M$.

Observation 2: As noted in Lemma 1, for $w \in V(P^*)$, $T' \notin w \Rightarrow T \in w$ and \exists a vertex $y \in V(P^*)$ s.t. $y = w \cup T' - T$.

Observation 3: P corresponds to $F_{P^*}(M')$.

Observation 4: (P^*, x^*, y^*) is a Dantzig figure. \therefore Since $\delta_{P^*}(x^*, y^*) = d^* (=d+m)$, \exists an NR path between x^* and y^* on P^* .

Notation

Let R^* be the NR path on P^* between x^* and y^* . $R^*: x^* = v_1^*, \dots, v_{d^*}^* = y^*$.

Let $B'_{v_i} = M' \cap v_i$, i.e., all primed symbols in v_i .

$$A'_{v_i} = M' - B'_{v_i}.$$

$$A_{v_i} = \{T \in M \mid T' \in v_i, T \notin v_i\}.$$

Observation 5:

$$\forall v_i \in V(P^*), v_i = y_i \cup B'_{v_i} \cup A_{v_i} \text{ where } y_i \in V(P).$$

Construction 2:

Form a sequence of vertices $R' : w_1, \dots, w_d^*$ s.t. $w_i = v_i \cup A'_{v_i} - A_{v_i}$,

where $v_i \in R^*$.

Lemma 2:

$$i) \quad w_1 = x \cup M' \in F_{P^*}^*(M').$$

$$ii) \quad w_i \in F_{P^*}^*(M') \quad \forall i.$$

$$iii) \quad w_d^* = y \cup M' \in F_{P^*}^*(M').$$

$$iv) \quad w_i, w_{i+1} \text{ are either neighbors or the same vertex, } \forall i.$$

$$v) \quad \text{If } w_i, w_{i+1} \text{ are the same vertex, delete one of them from } R'.$$

Then R' is an NR path from $x \cup M'$ to $y \cup M'$ on $F_{P^*}^*(M')$.

Proof:

$$i) \quad w_1 = v_1 \cup \phi - \phi = x^* = x \cup M' \in F_{P^*}^*(M').$$

$$\begin{aligned} ii) \quad w_j &= v_i \cup A'_{v_i} - A_{v_i} \\ &= \left(y_i \cup B'_{v_i} \cup A_{v_i} \right) \cup A'_{v_i} - A_{v_i} \quad (\text{by Observation 5}) \\ &= y_i \cup B'_{v_i} \cup A'_{v_i} = y_i \cup M'. \quad \therefore w_i \in F_{P^*}^*(M'). \end{aligned}$$

$$\begin{aligned}
\text{iii) } w_d^* &= v_d^* \cup A_{v_d^*}' - A_{v_d^*} \\
&= y^* \cup A_y' - A_y \\
&= (y \cup M) \cup M' - M \\
&= y \cup M' \in F_{P^*}(M').
\end{aligned}$$

iv) Consider the transition from v_i to v_{i+1} on R^* . There are three possible cases.

Case 1: $v_{i+1} = v_i \cup S - S'$ where $S' \in M'$, $S \in M$

$$\begin{aligned}
\therefore w_{i+1} &= v_{i+1} \cup A_{v_{i+1}}' - A_{v_{i+1}} \\
&= v_{i+1} \cup (A_{v_i}' \cup S') - (A_{v_i} \cup S) \\
&= (v_{i+1} \cup S' - S) \cup A_{v_i}' - A_{v_i} \\
&= v_i \cup A_{v_i}' - A_{v_i} \\
&= w_i.
\end{aligned}$$

$$\therefore w_{i+1} = w_i.$$

Case 2: $v_{i+1} = v_i \cup S - T$ where $S \in P$, $T \in P$

$$\begin{aligned}
\therefore w_{i+1} &= v_{i+1} \cup A_{v_{i+1}}' - A_{v_{i+1}} \\
&= (v_i \cup S - T) \cup A_{v_i}' - A_{v_i} \\
&= (v_i \cup A_{v_i}' - A_{v_i}) \cup S - T \\
&= w_i \cup S - T.
\end{aligned}$$

$\therefore w_i$ and w_{i+1} are neighbors.

Case 3: $v_{i+1} = v_i \cup S - T'$ where $S \in P$, $T' \in M'$

$$\begin{aligned}
 \therefore w_{i+1} &= v_{i+1} \cup A'_{v_{i+1}} - A_{v_{i+1}} \\
 &= (v_i \cup S - T') \cup (A'_{v_i} \cup T') - (A_{v_i} \cup T) \\
 &= (v_i \cup A'_{v_i} - A_{v_i}) \cup S - T \\
 &= w_i \cup S - T.
 \end{aligned}$$

$\therefore w_i$ and w_{i+1} are neighbors.

Since R^* is an NR path on P^* , these are the only three possible cases.

v) Immediate from i - iv). ■

Corollary 2:

\exists an NR path R from x to y on P .


Proof:

Immediate from Lemma 2, Part V, and Observation 3. ■

4. APPENDIX


Example of Construction 1

Suppose we have a Polytope P' as follows:




and want NR path from x to y .


Eliminate symbols in common to form face P



Iteration 1: Choose symbol F



Iteration 2: Choose symbol c



Below are examples of paths R^* from x^* to y^* on P^* and corresponding R from x to y on P according to Construction 2.

- (1) R^* : $ABC'F' - ABC'F - BCC'F - CC'DF - CDEF$
 R : $AB \text{ --- } BC \text{ --- } CD \text{ --- } DE$
- (2) R^* : $ABC'F' - ABCF' - ACFF' - CEFF' - CDEF$
 R : $AB \text{ --- } AF \text{ --- } EF \text{ --- } DE$
- (3) R^* : $ABC'F' - BCC'F' - CC'DF' - CC'DF - CDEF$
 R : $AB \text{ --- } BC \text{ --- } CD \text{ --- } DE$
- (4) R^* : $ABC'F' - AC'FF' - C'EFF' - C'DEF - CDEF$
 R : $AB \text{ --- } AF \text{ --- } EF \text{ --- } DE$
- (5) R^* : $ABC'F' - BCC'F' - BCC'F - CC'DF - CDEF$
 R : $AB \text{ --- } BC \text{ --- } CD \text{ --- } DE$
- (6) R^* : $ABC'F' - AC'FF' - ACFF' - CEFF' - CDEF$
 R : $AB \text{ --- } AF \text{ --- } EF \text{ --- } DE$

References

- [1] Adler, I., "Abstract Polytopes," Ph.D. Thesis, Department of Operations Research, Stanford University, Stanford, California, (1971).
- [2] Klee, V. and D. W. Walkup, "The d-Stop Conjecture for Polyhedra of Dimension $d < 6$," Acta Mathematica, 117, (1967), 53-78.